



## HEALTH & HYGIENE

The consumer requires meat that is free of harmful bacteria and drug residues. Special care should be taken to observe withdrawal times for health treatments.

Effective cleaning and length of time between flocks are important in the prevention of infection from one flock of broilers to the next. Usually a house should be rested no less than two weeks between flocks.

Dismantle and remove all equipment from the house. Thoroughly clean and disinfect the walls and floors. Drain the waters as well as the water storage tank. Remove soiled discarded litter from the vicinity of the house where it can be buried, burnt, or composted for fertilizer.

### Density Guidelines

Birds per sq. ft.....1 to1.2  
 Waterers (bells).....1 per 100 birds  
 Waterers (trough).....1 in. per bird  
 Waterers (nipples).....8 to 10 per nipple  
 Feeders (pans).....30 to 50 birds / pan  
 Feeders (trough).....2 in. per bird

## WATER MEDICATIONS FOR DISEASE CONTROL

Coccidiosis.....Amprolium  
 Chronic Respiratory.....Tylosin  
   Norfloxacin  
 Fowl Cholera.....Tetracyclines  
   Norfloxacin  
 Coryza.....Erythromycin  
   Sulfadimethoxine  
 Colibacillosis.....Neomycim  
   Norfloxacin  
 Enteritis..... Bacitracin  
   Lincomycin  
 Crop Mycosis.....Copper Sulfate  
   Iodine  
 Worms.....Piperazine  
   Flubendazole

## WATER ADDITIVE DOSAGES

Sugar.....1/4 lb/gal  
 Vinegar.....1qts/50gal  
 Chlorine Bleach.....1/2 cup/100gal  
 Skim Milk Powder.....1/4 lb/10gal  
 Aspirin.....85 gm/liter

## BASIC LIGHTING PRO- GRAM

Day 1 to 7.....23 hrs light  
 Day 8 to 28.....No light at night  
 Day 29 to market.....23 hrs light



## **BROILER FEEDING & MANAGEMENT**



## Reimers Feed Mill

**Spanish Lookout**

**Belize City, Belize, C.A.**

**Tel: (501) 823-0105**

**Fax: (501) 823-0229**

**Email: [sales@reimersfeed.com](mailto:sales@reimersfeed.com)**